

UNIVERSITY OF WATERLOO

Department of Systems Design Engineering

SD311

Engineering Optimization

Spring 2002

INSTRUCTOR: Paul H. Calamai, DC-2623 (x3182) , phcalamai@solstice

Midterm Exam : Thursday June 13, 2002

NAME:

ID:

PLEASE NOTE:

1. There are 8 problems to complete in order to obtain full marks.
2. The time limit for the examination is 2 hours.
3. Read over all of the questions before starting ... you will find that some problems are easier than others. The marking scheme is outlined in the chart below.
4. In addition to this title page there should be 10 pages to this exam. Pages 6 and 10 are intentionally blank.
5. Drawing instruments are the only aids permitted.

GOOD LUCK

#1	10	
#2	10	
#3	10	
#4	10	
#5	15	
#6	10	
#7	15	
#8	20	
Total	100	

1. (Total 10%) A furniture manufacturer has 3600 board feet of Walnut, 4300 board feet of Maple, and 6550 board feet of Oak in stock. The manufacturer can produce three products using this lumber, with input requirements as given below:

	Wood needed (bd. ft./unit)			Profit
	Walnut	Maple	Oak	\$/unit
Table	10	50	100	100
Desk	10	30	40	50
Chair	80	5	0	10

A table is always sold in combination with 4 chairs; and a desk is always sold in combination with one chair. However, a chair can be sold independently. Formulate an LP model to determine the product mix that maximizes profit. Be sure to state any assumptions that you have to make!

2. (Total 10%) Solve the following linear program graphically. Be sure to clearly label constraints boundaries and objective function contours as well as the feasible region. Also indicate what solution(s) is(are) optimal.

$$\begin{array}{llllll} \text{Minimize} & -y_1 & + & 2y_2 & & \\ \text{subject to} & y_1 & - & y_2 & \geq & 5 \\ & & & -y_2 & \leq & 10 \\ & 2y_1 & + & y_2 & \leq & 20 \\ & y_1 \geq 0, y_2 \leq 0 & & & & \end{array}$$

3. (Total 10%) Write the following problem in LP standard form as a **maximization** problem:

$$\begin{array}{ll}\text{Minimize} & -53x_1 + 33(x_1 - \frac{1}{3}x_3) \\ \text{subject to} & x_j + 1 \leq x_{j+1} \qquad j = 1, 2 \\ & x_1 + x_2 = 10 \\ & x_1 \leq 0, x_3 \geq 0\end{array}$$

4. (Total 10%) Consider the following LP standard display:

	x_1	x_2	x_3	x_4	
max c	10	1	0	0	b
	-1	1	4	21	11
	2	6	0	-2	2
	B	N	B	N	

(a) Compute the current basic solution.

(b) Compute all simplex directions at the current basic solution.

(c) Verify that the simplex directions at the current basic solution are feasible.

(d) Determine whether each of the simplex directions is improving or not.

(e) For any improving simplex direction at the current basic solution determine the corresponding maximum feasible step and the new basis that would result after such a step in that direction.

5. (Total 15%) Consider the following LP:

$$\begin{array}{rcccccccl} \text{Maximize} & 10x_1 & + & 4x_2 & + & 3x_3 & & & \\ \text{subject to} & x_1 & & & & - & x_3 & \leq & 0 \\ & -x_1 & + & x_2 & + & 3x_3 & & \leq & 4 \\ & x_1 & + & x_2 & & & & \leq & 3 \\ & & & & & & x_1, x_2, x_3 & \geq & 0 \end{array}$$

Begin by (a) putting the above problem into LP Standard Form. Then (b) form the initial **dictionary** corresponding to this LP. Next (c) perform, via dictionaries, two complete pivots (i.e. iterations) of the simplex method as described in your handouts (including all the corresponding information). Finally (d) interpret the solution in your final dictionary.

(Note: Please be careful – if you start with an incorrect initial dictionary you can receive only half marks at best!)

6. (Total 10%) Solve the following two linear programs:

(a)

$$\begin{array}{llllll}
 \text{Minimize} & 3x_1 & + & 5x_2 & & \\
 \text{subject to} & x_1 & + & x_2 & \geq & 50 \\
 & x_1 & + & x_2 & \leq & 95 \\
 & & & x_2 & \leq & 30 \\
 & x_1 & & & \geq & 0
 \end{array}$$

(b)

$$\begin{array}{llllll}
 \text{Maximize} & 450v & + & 200c & & \\
 \text{subject to} & 10v & + & 7c & \leq & 70000 \\
 & v & + & c & = & 10000 \\
 & v & & & \leq & 7000 \\
 & & & c & \leq & 7000
 \end{array}$$

Multiple Choice problems, 6@2% + 1@3% = Total 15%

7. In each of the following circle whichever choice provides the best answer.

The first two questions relate to the following standard form linear program Model 1:

$$\begin{array}{llllll} \text{Minimize} & 3x_1 & + & 2x_2 & + & 9x_3 \\ \text{subject to} & x_1 & + & x_2 & + & x_3 & = & 9 \\ & x_1 & - & x_2 & & & = & 4 \\ & x_1 & , & x_2 & , & x_3 & \geq & 0 \end{array}$$

- (a) Which of the following is an extreme point solution to Model 1?
- (i) $x = (3, 6, 0)$ (iii) $x = (4, 0, 5)$
(ii) $x = (3, 3, 3)$ (iv) $x = (6, 2, 1)$
- (b) Which of the following is **not** required of a feasible direction at $x = (6.5, 2.5, 0)$ in Model 1?
- (i) $\Delta x_1 + \Delta x_2 + \Delta x_3 = 0$ (iii) $\Delta x_1 \geq 0$
(ii) $\Delta x_1 - \Delta x_2 = 0$ (iv) $\Delta x_3 \geq 0$

The next two questions relate to the following maximizing standard form linear program Model 2:

	w_1	w_2	w_3	w_4	w_5	
max c	-1	3	0	7	-2	b
A	2	-1	1	4	1	8
	0	1	3	0	3	12

- (c) If variables w_1 and w_3 are currently basic in Model 2, the simplex direction for w_5 in the next move in Simplex solution of Model 2 is:
- (i) $\Delta w = (0, 0, 1, 0, 1)$ (iii) $\Delta w = (2, 0, 4, 0, 1)$
(ii) $\Delta w = (0, 0, -1, 0, 1)$ (iv) $\Delta w = (0, 0, -1, 0, 0)$
- (d) The maximum feasible step α for a move in the direction $\Delta w = (-6, -1, 1, 0, 0)$ from solution $w = (6, 6, 1, 0, 1)$ of Model 2 is:
- (i) 4 (iii) 2
(ii) 3 (iv) 1

The final three Multiple Choice questions are:

- (e) If the objective function of a general mathematical program is unimodal then every local optimum is a global optimum.
- (i) True (ii) False
- (f) If the main (i.e. functional) constraints of a general mathematical program are linear then its feasible set is necessarily convex.
- (i) True (ii) False
- (g) (3%) Suppose the two-phase improving search of a general mathematical program with original variables x_1, x_2 , and x_3 terminates with the following Phase 1 solutions in different cases. Identify all cases that would proceed with Phase II.
- (i) Global optimum $x = (40, 7, 0, 9, 0)$. (iii) Local optimum $x = (1, 3, 1, 0, 0)$
(ii) Global optimum $x = (22, 4, 3, 0, 0)$ (iv) Local optimum $x = (0, 2, 6, 5, 1)$, which may not be a global optimum.

8. (Total 20%) A refinery takes three raw gasolines and blends them to produce two types of fuel. Here are the data:

Raw gas type	Octane rating (Oc.R)	Supply (barrels/day)	Cost (\$/barrel)
1	68	4000	31.02
2	86	5050	33.15
3	91	7100	36.35

Fuel type	Minimum Oc.R	Profit (\$/barrel)	Demand (barrels/day)
1	95	45.15	At most 10,000
2	85	40.99	At least 15,000

The company sells raw gasoline not used in making fuels at \$38.95/barrel if its octane rating is greater than 90, and at \$36.85/barrel otherwise.

Formulate an LP model to determine the compositions of the two fuels to maximize total daily profit. Be sure to state any assumptions that you've made!

